



SOP

Collection of peer reviewed
scientific papers

-Agro sector-

Index:

Reducing N Fertilization without yield penalties in maize with a commercially available seed dressing (SOP COCUS MAIZE+).	
Published in "Agronomy", 24/02/2021	3
A new phytostimulating treatment for maize seed germination.	
17 th IFOAM Organic World Congress (Namyangju City, Rep. of Korea), 2011	5

Reducing N Fertilization without Yield Penalties in Maize with a Commercially Available Seed Dressing (SOP COCUS MAIZE+)

V. Tabaglio¹, A. Fiorini¹, M. Chiodini^{2,3}, L. Poggianella⁴

¹Department of Sustainable Crop Production (DI.PRO.VE.S.), Università Cattolica del Sacro Cuore, Piacenza, Italy;

²Università degli Studi di Milano, Italy; ³SOP Srl., Busto Arsizio, Italy; ⁴UC Davis, USA

The Background

Conventional management of agricultural production has often depleted soil quality and decreased soil health. Depleted soils can further reduce the provision of essential ecosystem services for crop production and local environments, impacting yields and water use efficiency.

The combination of intensive tillage and high nitrogen (N) fertilization also increases the mineralization of soil organic carbon (SOC), a process that releases CO₂ emissions and exacerbates the contribution of agricultural soils to Greenhouse Gas concentrations (GHGs) in the atmosphere [1]. Finding solutions that work within farmer economics to maintain crop production while mitigating environmental and climate impacts is an essential step toward a more sustainable world.


The Study

The study (<https://doi.org/10.3390/agronomy11030407>) was designed to evaluate the possibility to reduce synthetic N fertilization in the cultivation of corn (*Zea mays* L.). The trial was conducted at CERZOO research center (Piacenza, Northern Italy), under real world field conditions using existing agricultural soil, in 200 m² plots, randomized to minimize the influence of any potential soil differences upon the final results.

Three levels of fertilization were tested: 230 kg (100%), 160 kg (-30%) and 0 kg of synthetic N were applied in different plots after a common fertilization with 50 kg of equivalent efficient N from cattle slurry, with SOP COCUS MAIZE+ and without (Control). The roots were measured at anthesis and the crop cycle N₂O emissions were measured for the entire crop cycle year (April 2019 - April 2020).

The Results

A comparison analysis was conducted for root structure, crop yield, and soil N₂O emissions between the Control plots using fertilization at 100% synthetic N and the SOP COCUS MAIZE+ plots using -30% less synthetic N. Field data demonstrated a significant increase for the SOP treated plants in the very fine root hairs and the fine root length, both of which improve the efficiency of water uptake and plant nutrient cycling. Yield remained stable for both plots, Control and SOP (approximately 14 tons of grain per hectare), while N₂O emissions decreased by -23% per unit of production in the SOP treated plots.



SOP & the UN Sustainable Development Goals (SDGs)

The possibility to reduce synthetic N fertilization while maintaining crop yields would represent a step forward toward both environmental sustainability, human health, and overall food security under increasing risks from climate change. The crucial link between the UN Sustainable Development Goals (SDGs) and agricultural production is based upon the efficient use of nitrogen in cereal production systems [2]. The European Commission (EU) has also recognized this crucial link, and has set ambitious goals for reducing synthetic fertilizer usage by -20% at the field level by 2030 [3]. The primary reasons for these regulatory guidelines are the significant impacts upon the environment and human health from the inefficient use synthetic fertilization in agricultural production [4]. Most importantly, it is generally known that the excess use of synthetic N fertilization does not necessarily lead to increased crop yields, but can instead lead to high N losses in surface- and groundwater bodies causing negative water quality impacts, in addition to increased atmospheric GHGs emissions via N₂O cycling [5].

SOP products demonstrate the ability to improve the environmental footprint of agricultural operations and to reduce the risks associated with climate change for staple crop production. By establishing a commercial and scientific basis for maintaining current crop production levels with reduced need for synthetic fertilization, SOP products support the UN SDG framework for a more sustainable world and can be recognized by producers and their supply chains for SDG6 (Water Quality), SDG11 (Sustainable Cities), SDG12 (Responsible Production), SDG13 (Climate Action), and SDG15 (Life on land). The SOP COCUS Maize+ application has been certified according to ISO standards and the ISO warrants the recognition of alignment with the UN SDG goals with this SOP application in commercial use.

The Research Team

Prof. Dr. Vincenzo Tabaglio and Dr. Andrea Fiorini work in the department of Sustainable Crop Production (DI.PRO.VE.S.) at Università Cattolica del Sacro Cuore, located in Piacenza, Italy. Their work focuses on the study of new practices and innovative technologies for crop production intended for animal feedstock and food production, with a view toward sustainability within the food industry. Dr. Marcello Chiodini, PhD, a researcher at the University of Milan and a SOP collaborator, and Lorenzo Poggianella, from the University of California, Davis, assisted with the crop trial design and the analysis of the data sets.

The SOP Product

SOP COCUS MAIZE+ is a seed dressing based on natural materials and is produced by SOP - Save Our Planet. The SOP technical process creates crop applications that work with the microbial communities present in the soil and the rhizosphere in order to improve plant-microbial interaction. The goal of the SOP COCUS Line is to improve water use efficiency, plant nutrient availability, and nutrient uptake in corn production for increased plant resilience and improved farmer economics. This application has been commercialized since 2005 and is used by farmers in several EU countries.

[1] Jones, A.; Panagos, P.; Barcelo, S.; Bouraoui, F.; Bosco, C.; Dewitte, O.; Gardi, C.; Erhard, M.; Hervas, J.; Hiederer, R. et al. *The State of Soil in Europe: A Contribution of the JRC to the European Environment Agency's Environment State and Outlook Report—SOER 2010*; European Commission: Luxembourg, 2012, doi:10.2788/77361.

[2] Ladha, J.K.; Jat, M.L.; Stirling, C.M.; Chakraborty, D.; Pradhan, P.; Krupnik, T.J.; Sapkota, T.B.; Pathak, H.; Rana, D.S.; Tesfaye, K.; et al. Chapter Two – Achieving the sustainable development goals in agriculture: The crucial role of nitrogen in cereal – based systems. In *Advances in Agronomy*; Sparks, D.L., Ed; Academic Press: Cambridge, MA, USA, 2020; Volume 163, pp. 39–116.

[3] <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=COM:2020:0381:FIN>.

[4] https://www.organicseurope.bio/content/uploads/2020/06/2016_RISE_NRR_Full_EN_compressed.pdf?dd

[5] Good, A.G.; Beatty, P.H. Fertilizing Nature: A Tragedy of Excess in the Commons. *PLoS Biol.* 2011, 9, e1001124, doi:10.1371/journal.pbio.1001124.

Presented at:
17th IFOAM
Organic World Congress
(Namyangju City, Rep. of Korea - 2011)

A new phytostimulating treatment for maize seed germination

A. Zanierato¹, D. Tassi², I. Accorinti¹, P. Luparia¹, E. Vieira de Manincor³, C. Baffi⁴

¹SOP Srl, Busto Arsizio (VA), Italy; ²Vittorio Tadini Experimental Farm, Podenzano (PC), Italy; ³Independent Professional Agronomist, Italy; ⁴Istituto di Chimica Agraria ed Ambientale, Faculty of Agriculture, Piacenza (PC), Italy


Objectives

The aim of this study was to evaluate the efficacy of a bio-enhancer (SOP), to be applied as a seed dressing before sowing, on the germination speed of two varieties of Maize caryopses, belonging to two different classes of precocity, on an experimental farm in northern Italy.

Materials & Methods

Formula	SOP SQG 377
Seed class	Class 400 hybrid LG 33.9, class 700 hybrid LG AZUAGA
Materials & Methods	Two varieties of maize belonging to two different classes of precocity were compared. Undressed seeds were dressed differently before they were planted in pots. The samples for each class were: 1) undressed control; 2) dressed with SQG 377 200 g/ha (recommended standard dosage); 3) dressed with conventional fungicide at recommended dressing dosage.
Evaluated parameters	Seed emergence
Statistical significance	P<0.05

Results & Graphs



Conclusions

SOP helps enhance emergence speed and uniformity of maize caryopses, giving advantages to both early and late sowing.